

California State University, Monterey Bay

Digital Commons @ CSUMB

CSU Student Research Competition Delegate
Entries

Undergraduate Research Opportunities Center
(UROC)

4-29-2017

Examining Temperament in Exercise Dependence and Eating Disorders

James Vongsaroj

California State University, Monterey Bay

Follow this and additional works at: https://digitalcommons.csumb.edu/uroc_csusrc

Recommended Citation

Vongsaroj, James, "Examining Temperament in Exercise Dependence and Eating Disorders" (2017). *CSU Student Research Competition Delegate Entries*. 2.
https://digitalcommons.csumb.edu/uroc_csusrc/2

This Presentation is brought to you for free and open access by the Undergraduate Research Opportunities Center (UROC) at Digital Commons @ CSUMB. It has been accepted for inclusion in CSU Student Research Competition Delegate Entries by an authorized administrator of Digital Commons @ CSUMB. For more information, please contact digitalcommons@csumb.edu.

Examining Temperament in Exercise Dependence and Eating Disorders

James Vongsaroj

Faculty Mentor: Brian Cook, Ph.D.
Kinesiology Department, CSUMB

California State University Research Competition
San Luis Obispo, CA
April 29, 2017



California State University
MONTEREY BAY
Extraordinary Opportunity

Background

- Over exercise can have serious implications
- 25% - 55% in patients with eating disorders
- Health may be impaired in at-risk individuals
 - 20% - 40% mortality rate – highest of all mental illness
 - Over exercise is implicated in this
- Why do some people exercise too much?



Background

- Exercise Dependence (EXD) is defined as when exercise results in physical, psychological, and/or social detriments

Two distinct Variants:

- Primary Exercise Dependence
 - Exercise is the only problem
- Secondary Exercise Dependence
 - Exercise to facilitate other disorders
 - Most often - eating disorders



Background

- Temperament is an individual's predisposition of self-regulation processes
 - Behavioral Activation System (**BA**)
 - impulsivity and seeking rewarding consequences
 - Behavioral Inhibition System (**BI**)
 - sensitivity to punishment, non-reward, novelty, anxiety, and depression



Background & Gap

- Recent research has found
 - **BA** in elite athletes with EXD
 - **BI** in hospitalized inpatients with eating disorders

My Research Question

Which temperament styles are associated with each EXD variant and eating disorders risk in a general-population based sample?



Purpose & Hypotheses

- Purpose:
 - To examine temperament styles in EXD and eating disorders risk in a general-population based sample
- Hypotheses:
 - Individuals with eating disorders risk will be associated with **BI** temperament
 - Individuals with EXD will be associated with **BA** temperament



Participants

- 880 individuals who completed an online survey
 - Mean age = 28.46+10.13
 - Mean Body Mass Index = 27.29+6.34
 - 63.92% female
 - 66.90% Caucasian
 - 42.41% never married/currently single
 - 29.60% earned a college diploma
 - 60.41% with an annual personal income of <\$25k
 - 79.63% were not Hispanic or Latino



Measures

- Demographics Questionnaire
- Exercise Dependence Scale (EDS)
 - 21 Items measured on a 6-point Likert scale
 - Reliability in this study was excellent ($\alpha = 0.95$)
- Eating Disorders Examination-Questionnaire (EDE-Q)
 - 28-item questionnaire evaluating symptoms of an eating disorder
 - Continuous symptom score calculated to determine risk
 - Reliability in this study was excellent ($\alpha = 0.90$)



Measures

- Leisure-Time Exercise Questionnaire (LTEQ):
 - Self-report questionnaire that assesses 20 minute bouts of strenuous, moderate, and light intensity exercise
- Behavioral Inhibition/Behavior Activation Scales (BIS/BAS):
 - 24 item scale assessing BI & BA
 - BAS subscales – Drive, Fun Seeking, & Reward Responsiveness
 - Reliability in this study was good for the BIS ($\alpha = 0.81$) and BAS ($\alpha = 0.86$)



Procedures

- Reviewed and approved by IRB
- Informed consent collected prior to completing an online survey
- Participants were grouped as:
 - Regular exercisers (e.g., LTEQ > 24.00)
 - Primary EXD (e.g., EDS > 77.00 & EDE-Q < 2.98)
 - Secondary EXD (e.g., EDS > 77.00 & EDE-Q > 2.99)
 - Eating disorders risk only (e.g., EDS < 77.00 & EDE-Q > 2.99)



Statistical Analyses

One-Way ANOVA

- Overall group differences
 - Tukey Post Hoc analyses were conducted to examine individual group differences
- Dependent variables – BIS & BAS scores (continuous)
- Independent variables – Groups based on LTEQ, EDS, & EDE-Q scores (categorical)



ANOVA Results

	Regular Exercisers M(SD)	Primary Exercise Dependence M(SD)	Secondary Exercise Dependence M(SD)	Eating Disorders Risk Only M(SD)		<u>ANOVA</u>	
					df	F	η^2
BIS	19.21 (4.16)	19.40 (4.25)	18.83 (4.01)	21.90 (4.15)	3, 841	27.13*	.088
BAS	39.27 (6.00)	39.62 (8.37)	37.25 (7.67)	39.50 (6.41)	3, 830	1.52	.005
BAS-Drive	11.16 (2.50)	11.70 (2.97)	10.79 (2.35)	11.02 (2.72)	3, 850	0.87	.003
BAS-Fun	11.44 (2.40)	11.56 (2.74)	11.20 (3.01)	11.67 (2.56)	3, 852	0.78	.003
BAS-Reward	16.68 (2.57)	16.35 (3.67)	15.20 (3.61)	16.89 (2.74)	3, 850	4.76*	.017

Table 1: Groups means for study outcomes and ANOVA results.

NOTES: BIS = Behavioral Inhibition Scale; BAS = Behavioral Activation Scale; M(SD) = Mean (Standard Deviation); df = degrees of freedom; * = $p < .001$; η^2 = effect size.



Post Hoc Analyses Results




Figure 1: Group differences for Behavioral Inhibition and Behavioral Activation Scales

NOTE: BIS = Behavioral Inhibition Scale; BAS = Behavioral Activation Scale;

* = $p < .05$; ** = $p < .01$



Discussion

- Key Findings!!
- Hypothesis 1 – BI & eating disorders risk
 - **Supported!**
 - **Eating disorders risk only group reported highest scores in overall BIS**
- Hypothesis 2 –BA & exercise dependence
 - **Partially Supported!**
 - **BAS Reward subscale only showed significant differences**



Discussion

- General-population sample
- Provides new insights on temperament styles
 - Secondary EXD lowest BAS Reward
 - Exercise & eating disorder risk – may not be rewarding
- Temperament styles may be more heterogeneous than previously thought
 - More research needed on reward systems



California State University
MONTEREY BAY
Extraordinary Opportunity

Limitations

- Cross-sectional design
 - Can't prove cause and effect
- Self-report measures - limits accuracy of assessments
 - No clinical diagnosis
 - EXD or eating disorders
 - No objective assessments of physical activity



Conclusions

- BAS may be more complex than previously reported
- Clinical implications
 - A person's temperament may explain why they exercise
 - Tailor interventions to match temperament style




References

- American Psychiatric Association Diagnostic and Statistical Manual of Mental Disorders (DSM-IV-TR). (2000) Washington, DC.
- Bamber, D., Cockerill, I., Rodgers, S., Carroll, D. (2003). Diagnostic criteria for exercise dependence in women. *British Journal of Sports Medicine*, 37(5),393-400.
- Carver S., White T. (1994) "Behavioral Inhibition, Behavioral Activation, and Affective Responses to Impending Reward and Punishment: The BIS/BAS Scales." *Journal of Personality and Social Psychology* 67(2), 319-33.
- Cook, B., Hausenblas, H., & Freimuth, M. (2014). Exercise addiction and compulsive exercising: Relationship to eating disorders, substance use disorders and addictive disorders. *Eating disorders, addictions and substance use disorders*, 127-144.
- Derryberry, D., & Rothbart, M. K. (1984). Emotion, attention, and temperament. *Emotions, cognition, and behavior*, 132-166.
- Fairburn, C. G., & Beglin, S. J. (2008). Eating Disorder Examination Questionnaire (EDE-Q 6.0). In C. G. Fairburn (Ed.), *Cognitive behavior therapy and eating disorders*, 309–314.
- Godin, G., & Shepard, R.J. (1985). A simple method to assess exercise behavior in the community. *Canadian Journal of Applied Sports Science*, 10, 141-146.
- Hausenblas, H., Symons D. (2002). Exercise dependence: a systematic review. *Psychology of Sport and Exercise*, 3(2), 89–123.
- Meyer, C., Taranis, L., (2011). Exercise in the eating disorders: Terms and definitions. *European Eating Disorders Review*, 19(3), 169-17.
- Müller, A., Claes, L., Wos, K., Kerling, A., Wünsch-Leiteritz, W., Cook, B., & de Zwaan, M. (2015). Temperament and Risk for Exercise Dependence: Results of a Pilot Study in Female Patients with Eating Disorders Compared to Elite Athletes *Psychopathology*, 48(4), 264-269.
- Peterson, C. B., Crosby, R. D., Wonderlich, S. A., Joiner, T., Crow, S. J., Mitchell, J. E., & Le Grange, D. (2007). Psychometric properties of the eating disorder examination questionnaire: Factor structure and internal consistency. *International Journal of Eating Disorders*, 40(4), 386-389.
- Shroff, H., Reba, L., Thornton, M.L., Tozzi, F., Klump, K.L., Berrettini, W.H...& Bulik, C. (2006). Features associated with excessive exercise in women with eating disorders. *International Journal of Eating Disorders*, 39(6), 454-461.



Acknowledgements

- The CSU Research Competition
- CSUMB Undergraduate Research Opportunities Center
- CSUMB Kinesiology Department
- Dr. Cook for his mentorship
- ***You for being a great audience!***



Questions?



California State University
MONTEREY BAY
Extraordinary Opportunity